

YITP long-term and Nishinomiya-Yukawa memorial workshop

Hadrons and Hadron Interactions in QCD 2024 (HHIQCD 2024)

-- Experiments, Effective theories, and Lattice --

14th Oct. - 15th Nov., 2024

Yukawa Institute for Theoretical Physics, Kyoto University, Japan



电子科技大学

University of Electronic Science and Technology of China

Effective range expansion with the left-hand cut

Meng-Lin Du (UESTC)

University of Electronic Science and Technology of China


In collaboration with A. Fillin, V. Baru, X.-K. Dong, E. Epelbaum, F.-K. Guo, C. Hanhart, A. Nefediev, J. Nieves, Q. Wang, and B. Wu

Based on PRL 131,131903 (2023), and arXiv:2408.09375[hep-ph]

Oct. 18, 2024 @ Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto, Japan

Hadrons and Hadron Interactions in QCD 2024 (HHIQCD2024)

Doubly charmed tetraquark (Tcc)




Breit-Wigner fit

LHCb, Nature Phys. 18, (2022) 751

Parameter	Value
N	117 ± 16
δm_{BW}	$-273 \pm 61 \text{ keV}$
Γ_{BW}	$410 \pm 165 \text{ keV}$

$\Rightarrow \mathfrak{R} \sim 400 \text{ keV}$.



Unitarized and analytical

LHCb, Nature Commun. 13 (2022), 3351

$$\delta m = m_{T_{cc}^+} - m_{D^{*+}} - m_{D^0}$$

$$\delta m_{\text{pole}} = -360 \pm 40_{-0}^{+4} \text{ keV}$$


$$\Gamma_{\text{pole}} = 48 \pm 2_{-14}^{+0} \text{ keV}$$

$\Rightarrow I = 0$: isoscalar

$\hookrightarrow D^+ D^0 \pi^+, D^+ D^+ \times$


$\Rightarrow T_{cc}^+$ resides near $D^* D$ thresholds

\hookrightarrow approximate 90% of $D^0 D^0 \pi^+$ events contain a D^{*+} .



LHCb, Nature Commun. 13 (2022)

The three-body cut



☞ Three-body cuts



☞ LO Chiral Lagrangian (g determined from $D^* \rightarrow D\pi$)



$$\mathcal{L} = \frac{1}{4} g \text{Tr} (\vec{\sigma} \cdot \vec{u}_{ab} H_b H_a^\dagger)$$

☞
$$U_\alpha(M, p) = P_\alpha - \sum_\beta \int \frac{d^3 \vec{q}}{(2\pi)^3} V_{\alpha\beta}(M, p, q) G_\beta(M, q) U_\beta(M, q)$$

$\hookrightarrow G_\alpha(M, p) = \frac{1}{m_\alpha^* + m_\alpha + \frac{p^2}{2\mu_\alpha} - M - \frac{i}{2} \Gamma_\alpha(M, p)}$

Description of the experimental data

Du et al., PRD 105, 014024(2022)



Scheme	III	II	I
Pole [keV]	$-356^{+39}_{-38} - i(28 \pm 1)$	$-333^{+41}_{-36} - i(18 \pm 1)$	$-368^{+43}_{-42} - i(37 \pm 0)$

Complete 3-bdoy cut

No OPE

No 3-body cut
But constant D^* width



👉 The width of T_{cc}^+

$$56 \text{ keV} \xrightarrow[\text{OPE}]{\text{remove}} 36 \text{ keV} \xrightarrow[\text{M-dep. of } \Gamma^*]{\text{remove}} 74 \text{ keV}$$

Doubly Charm Tetraquark on the Lattice


Padmanath *et al*, PRL129,032002(2022)

	m_D (MeV)	m_{D^*} (MeV)	M_{av} (MeV)	$a_{l=0}^{(J=1)}$ (fm)	$r_{l=0}^{(J=1)}$ (fm)	$\delta m_{T_{cc}}$ (MeV)	T_{cc}
Lattice ($m_\pi \simeq 280$ MeV, $m_c^{(h)}$)	1927(1)	2049(2)	3103(3)	1.04(29)	$0.96^{(+0.18)}_{(-0.20)}$	$-9.9^{+3.6}_{-7.2}$	Virtual bound st.
Lattice ($m_\pi \simeq 280$ MeV, $m_c^{(l)}$)	1762(1)	1898(2)	2820(3)	0.86(0.22)	$0.92^{(+0.17)}_{(-0.19)}$	$-15.0^{(+4.6)}_{(-9.3)}$	Virtual bound st.
Experiment [2,41]	1864.85(5)	2010.26(5)	3068.6(1)	-7.15(51)	$[-11.9(16.9), 0]$	-0.36(4)	Bound st.





$$t = \frac{E_{cm}}{2} \frac{1}{p \cot \delta - ip},$$

$$p \cot \delta = \frac{1}{a_0} + \frac{1}{2} r_0 p^2,$$





The three-body cut vs. left-hand cut






left-hand cut

$$\int_{-1}^1 d \cos \theta G_\pi(E, p, p)$$



$$G_\pi^{-1}(E, k, k') \xrightarrow[\text{on shell: } k=k'=p]{\cos \theta = \pm 1} E_{D^*}(p^2) - E_D(p^2) - \omega_\pi(4p^2/0) = 0$$

The left-hand cut



$$m_\pi = 280 \text{ MeV}$$

☞ two-body branch point:

$$E = M_D + M_{D^*}$$

$$\Rightarrow p_{\text{rhc}2}^2 = 0$$

☞ three-body branch point:

$$E = M_D + M_D + m_\pi$$

$$\Rightarrow \left(\frac{p_{\text{rhc}3}}{E_{DD^*}} \right)^2 = +0.019$$

☞ left-hand cut branch point:


$$\Rightarrow \left(\frac{p_{\text{lhc}}^{1\pi}}{E_{DD^*}} \right)^2 = -0.001$$

$$\left(\frac{\tilde{p}_{\text{lhc}}^{1\pi}}{E_{DD^*}} \right)^2 = -0.190$$

Phase shift with the left-hand cut: LSE


Du et al., PRL 131,131903 (2023)

$M_D = 1927 \text{ MeV}$, $M_{D^*} = 2049 \text{ MeV}$, $m_\pi = 280 \text{ MeV}$




Pole trajectory

$$M_D = 1927 \text{ MeV}, M_{D^*} = 2049 \text{ MeV}, m_\pi = 280 \text{ MeV}$$



bound state \longrightarrow virtual state \longrightarrow resonances below threshold



No virtual state along the $1hc$

Related recent works on FV w/ the lhc...


Also talked by Lyu on 14pm

- Plane-wave basis to treat long-range interactions
Project to irep. of the cubic to avoid the lhc associated to the partial wave projection
Meng and Epelbaum, JHEP (2021)
Meng et al., PRD (2024)
Mai and Döring, EPJA (2017), PRL (2019)
- Generalization of the Lüscher + K -matrix
Hansen and Raposo, JHEP (2024)
- Three-body framework (automatically includes lhc)
Dawid et al., PRD (2023)
Hansen et al., PRD (2024)
- Modify the Lüscher formula via “modified effective range expansion”
Bubna et al., JHEP (2024)

ERE

$$f(k^2) = \frac{1}{k \cot \delta - ik}$$

$$k \cot \delta = \frac{1}{a} + \frac{1}{2}rk^2 + \mathcal{O}(k^4)$$




The N/D method

$$T(s) = \frac{N(s)}{D(s)}$$

$$\text{Im}D = \text{Im}\frac{N}{T} = N\text{Im}\frac{1}{T} = \begin{cases} -N\rho, & s > s_{\text{thr}} \\ 0, & s < s_{\text{thr}} \end{cases}$$

$$\text{Im}N = \begin{cases} \text{Im}TD, & s < s_{\text{lhc}} \\ 0, & s > s_{\text{lhc}} \end{cases}$$



$$D(s) = \sum_i \frac{\gamma_i}{s - s_i} + \sum_{m=0}^{n-1} a_m s^m - \frac{(s - s_0)^n}{\pi} \int_{s_{\text{thr}}}^{\infty} ds' \frac{\rho(s')N(s')}{(s' - s)(s' - s_0)^n},$$


$$N(s) = \sum_{m=0}^{n-\ell-1} b_m s^m + \frac{(s - s_0)^{n-\ell}}{\pi} \int_{-\infty}^{s_{\text{left}}} ds' \frac{\text{Im}T(s')D(s')}{(s' - s_0)^{n-\ell}(s' - s)}.$$

N=1
→

$$D(s) = \sum_i \frac{\gamma_i}{s - s_i} + P(s) + G(s)$$

$$T(s) = \frac{1}{D(s)}$$

$$d(k^2) = P(k^2) - ik = \frac{1}{a} + \frac{1}{2}rk^2 + \mathcal{O}(k^4) - ik$$




Non-relativistic
Neglect CDD

The N/D method

$$T(s) = \frac{N(s)}{D(s)}$$


$$\text{Im}D = \text{Im}\frac{N}{T} = N\text{Im}\frac{1}{T} = \begin{cases} -N\rho, & s > s_{\text{thr}} \\ 0, & s < s_{\text{thr}} \end{cases}$$

$$\text{Im}N = \begin{cases} \text{Im}TD, & s < s_{\text{lhc}} \\ 0, & s > s_{\text{lhc}} \end{cases}$$



$$D(s) = \sum_i \frac{\gamma_i}{s - s_i} + \sum_{m=0}^{n-1} a_m s^m - \frac{(s - s_0)^n}{\pi} \int_{s_{\text{thr}}}^{\infty} ds' \frac{\rho(s')N(s')}{(s' - s)(s' - s_0)^n},$$

$$N(s) = \sum_{m=0}^{n-\ell-1} b_m s^m + \frac{(s - s_0)^{n-\ell}}{\pi} \int_{-\infty}^{s_{\text{left}}} ds' \frac{\text{Im}T(s')D(s')}{(s' - s_0)^{n-\ell}(s' - s)}.$$




$$\frac{1}{T_\ell^{\text{II}}} = \frac{1}{T_\ell} + 2i\rho \implies T^{\text{II}} = \frac{1}{\frac{D}{N} + 2i\rho} = \frac{N}{D + 2i\rho N}$$


Along the lhc, $i\rho$ and D is real, N has imaginary part.

$$D + 2i\rho N \neq 0$$

The left-hand cut arising from OPE



$$\text{Im } f(k^2) = c \text{Im } L(k^2) = -\frac{c}{4k^2} \pi, \quad \text{for } k^2 < k_{\text{thc}}^2$$



Solving LSE could be time-consuming.


For a t -channel exchange at low-energies, an S -wave amplitude reads

$$L_t(s) = \frac{1}{2} \int \frac{1}{t - m_5^2} d \cos \theta = -\frac{s}{\lambda(s, m_1^2, m_2^2)} \log \left(\frac{s - 2(m_1^2 + m_2^2) + m_5^2 + \frac{(m_1^2 - m_2^2)^2}{s}}{m_5^2} \right),$$

with m_5 the mass of changed particle. Likewise, the u -channel exchanged S -wave amplitude reads

$$L_u(s) = \frac{1}{2} \int \frac{1}{u - m_5^2} d \cos \theta = -\frac{s}{\lambda(s, m_1^2, m_2^2)} \left(\log(s + m_5^2 - 2(m_1 + m_2)^2) - \log(m_5^2 - \frac{(m_1^2 - m_2^2)^2}{s}) \right).$$

The left-hand cut: nonrelativistic



Exchanged-particle: relativistic

$$\eta = |m_1 - m_2| / (m_1 + m_2)$$

$$\mu_{\text{ex}}^2 = m_{\text{ex}}^2 - (m_1 - m_2)^2$$

$$\mu_+^2 = 4\mu\mu_{\text{ex}}^2 / (m_1 + m_2)$$

$$L_t(k^2) \equiv \frac{1}{2} \int_{-1}^{+1} \frac{d \cos \theta}{t - m_{\text{ex}}^2} = -\frac{1}{4k^2} \log \frac{m_{\text{ex}}^2/4 + k^2}{m_{\text{ex}}^2/4},$$

$$L_u(k^2) \equiv \frac{1}{2} \int_{-1}^{+1} \frac{d \cos \theta}{u - m_{\text{ex}}^2} \approx -\frac{1}{4k^2} \log \frac{\mu_+^2/4 + k^2}{\mu_+^2/4 + \eta^2 k^2},$$

$$\frac{1}{2} \int_{-1}^{+1} \frac{(\mathbf{p}_1 - \mathbf{p}_3)^2}{t - m_{\text{ex}}^2} d \cos \theta = -\frac{m_{\text{ex}}^2}{2} \int_{-1}^{+1} \frac{d \cos \theta}{t - m_{\text{ex}}^2} - 1$$

$$\frac{1}{2} \int_{-1}^{+1} \frac{(\mathbf{p}_1 - \mathbf{p}_4)^2}{u - m_{\text{ex}}^2} d \cos \theta \approx -\frac{\mu_{\text{ex}}^2}{2} \int_{-1}^{+1} \frac{d \cos \theta}{u - m_{\text{ex}}^2} - 1$$

$\mathcal{F}_l/2$

$$f(k^2) = \frac{n(k^2)}{d(k^2)}$$

$$\text{Im } d(k^2) = -k n(k^2), \quad \text{for } k^2 > 0,$$

$$\text{Im } n(k^2) = d(k^2) \text{Im } f(k^2), \quad \text{for } k^2 < k_{\text{lhc}}^2.$$

The N/D method: nonrelativistic

$$n(k^2) = n_m(k^2) + \frac{(k^2)^m}{\pi} \int_{-\infty}^{k_{\text{Ihc}}^2} \frac{d(k'^2) \text{Im} f(k'^2)}{(k'^2 - k^2)(k'^2)^m} dk'^2$$

$\propto \text{Im}L$

No singularity along Ihc

$$n(k^2) = n'_m(k^2) + \frac{P(k^2)}{\pi} \int_{-\infty}^{k_{\text{Ihc}}^2} \frac{\text{Im} f(k'^2)}{k'^2 - k^2} dk'^2 = n'_m(k^2) + P(k^2) \tilde{g}L(k^2)$$


$$\begin{aligned} n(k^2) &= n_0 + n_1 k^2 + \frac{k^2}{\pi} \int_{-\infty}^{k_{\text{Ihc}}^2} \frac{(d_0 + d_1 k'^2) \text{Im} f(k'^2)}{(k'^2 - k^2) k'^2} dk'^2 \\ &= n_0 + n_1 k^2 - cL_0 + (d_0 + d_1 k^2) \frac{c}{\pi} \int_{-\infty}^{k_{\text{Ihc}}^2} \frac{\text{Im} f(k'^2)}{k'^2 - k^2} dk'^2 \\ &= n'_0 + n_1 k^2 + (d_0 + d_1 k^2) cL(k^2) \end{aligned}$$

$$n(k^2) = \tilde{n}(k^2) + \tilde{g}(L(k^2) - L_0)$$

$$L_0 = L(k^2 = 0) = -1/\mu_{\text{ex}}^2$$

The N/D method: nonrelativistic

$$d(k^2) = d_n(k^2) - \frac{(k^2 - k_0^2)^n}{\pi} \int_0^\infty \frac{k' n(k'^2) dk'^2}{(k'^2 - k^2)(k'^2 - k_0^2)^n}$$



$$\begin{aligned} d(k^2) &= \tilde{d}(k^2) - ik(\tilde{n}(k^2) - \tilde{g}L_0) - \frac{\tilde{g}}{\pi} \int_0^\infty \frac{k' L(k'^2)}{k'^2 - k^2} dk'^2 \\ &= \tilde{d}(k^2) - ik n(k^2) - \tilde{g}d^R(k^2) \end{aligned}$$

$$d_u^R(k^2) = \frac{i}{4k} \left(\log \frac{\mu_+/2 + ik}{\mu_+/2 - ik} - \log \frac{\mu_+/2 + i\eta k}{\mu_+/2 - i\eta k} \right)$$

It is worth stressing that $d(k^2)$ is free of lhc, as the lhc associated with $n(k^2)$ below the threshold is counterbalanced by $d^R(k^2)$, which is crucial to ensure that $f(k^2)$ exhibits the correct lhc behavior. Along the rhc, both $n(k^2)$ and $d^R(k^2)$ are real such that $\text{Im}d(k^2) = -k n(k^2)$.

Effective range expansion with the left-hand cut

$$\frac{1}{f(k^2)} = \frac{\tilde{d}(k^2) - \tilde{g}d^R(k^2)}{\tilde{n}(k^2) + \tilde{g}(L(k^2) - L_0)} - ik$$

$$d_u^R(k^2) = \frac{i}{4k} \left(\log \frac{\mu_+/2 + ik}{\mu_+/2 - ik} - \log \frac{\mu_+/2 + i\eta k}{\mu_+/2 - i\eta k} \right)$$

$$L(k^2) = -\frac{1}{4k^2} \log \frac{\mu_+^2/4 + k^2}{\mu_+^2/4 + \eta^2 k^2}$$

$$\frac{1}{f_{[m,n]}(k^2)} = \frac{\sum_{i=0}^n \tilde{d}_i k^{2i} - \tilde{g}d^R(k^2)}{1 + \sum_{j=1}^m \tilde{n}_j k^{2j} + \tilde{g}(L(k^2) - L_0)} - ik$$

$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g}d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

$$\tilde{g} \rightarrow 0$$


$$\frac{1}{f(k^2)} = \frac{1}{a} + \frac{1}{2} r k^2 - ik$$

Scattering length $a = f(k^2 = 0) = \left[\tilde{d}_0 + \frac{\tilde{g}}{\mu_+} (1 - \eta) \right]^{-1}$

Effective range $r = \left. \frac{d^2(1/f+ik)}{dk^2} \right|_{k=0} = 2\tilde{d}_1 - \frac{8\tilde{g}}{3\mu_+^3} (1 - \eta^3) - \frac{4\tilde{g}}{\mu_+^4 a_u} (1 - \eta^4)$


Example: Tcc on the Lattice [3 parameters]

Du et al., 2408.09375 [hep-ph]




$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g} d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

Couplings to the exchanged-particle



$$g_P = -\frac{2\pi\tilde{g}}{\mu d^{0,\text{lhc}}\mathcal{F}_\ell}$$

$$g_{D^*D\pi}^2/(4F^2) = 9.2 \text{ GeV}^{-2}$$



$$-\frac{2\pi}{\mu}\text{Im}f = \text{Im}T = \text{Im}V_{\text{OPE}}(k^2) = g_P \frac{-\pi}{4k^2} \mathcal{F}_\ell, \quad \text{for } k^2 < k_{\text{lhc}}^2$$

$$\text{Im}n(k^2) = -\tilde{g} \frac{\pi}{4k^2}, \quad \text{for } k^2 < k_{\text{lhc}}^2$$

$$d_u^{0,\text{lhc}} = \tilde{d}_0 - \frac{\tilde{d}_1 \mu_+^2}{4} + \frac{\mu_+}{2} \left(1 + \frac{\tilde{g}}{\mu_{\text{ex}}^2} \right) + \frac{\tilde{g} \log[2/(1+\eta)]}{\mu_+}$$

The amplitude zero

At leading order, i.e., $\tilde{n}(k^2) = 1$,

$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g} d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

For a general u -channel exchange,

$$1 + \tilde{g} \left[L_u(k_{u,\text{zero}}^2) + \frac{1}{\mu_{\text{ex}}^2} \right] = 0,$$

for the case $|\Delta| \ll m_{\text{th}}$ such that $\eta \ll 1$,  the t -channel exchange

$$k_{t,\text{zero}}^2 = -\frac{m_{\text{ex}}^2}{4} \left[1 + \frac{1}{y} W(-e^{-y} y) \right]$$

where $y \equiv 1 + m_{\text{ex}}^2/\tilde{g}$ and W is the Lambert W function.

$$y = 1 + \frac{1 + \frac{4}{3} a_t m_{\text{ex}} (1 - \log 4) - \frac{4\pi a_t m_{\text{ex}}^2}{\mu_{gP} \mathcal{F}_\ell}}{2 + a_t m_{\text{ex}} (1 - m_{\text{ex}} r_t/4)}.$$

Summary

- The three-body cut: one-pion exchange + self-energy of D^*
- ★ Unphysical pion masses on the Lattice
 $M_D = 1927$ MeV, $M_{D^*} = 2049$ MeV, $m_\pi = 280$ MeV
↪ the three-body cut above the two-body cut ($\sqrt{s_{\text{thc}}} = 3968$ MeV)
↪ The traditional ERE valid only in a very limited range
↪ An accurate extraction of the pole requires the OPE implemented
- ★ The ERE with the left-hand cut

$$f_{[0,1]}(k^2) = \left[\frac{\tilde{d}_0 + \tilde{d}_1 k^2 - \tilde{g}d^R(k^2)}{1 + \tilde{g}(L(k^2) - L_0)} - ik \right]^{-1}$$

- ↪ correct behavior of the left-hand cut
- ↪ can be used to extract the couplings of the exchanged particle to the scattering particles
- ↪ amplitude zeros caused by the interplay between the short- and long-range interactions

Thank you very much for your attention!

Thank you very much for your attention!

Without $d^R(k^2)$

$$d(k^2) = \tilde{d}(k^2) - ik(\tilde{n}(k^2) - \tilde{g}L_0) - \frac{\tilde{g}}{\pi} \int_0^\infty \frac{k' L(k'^2)}{k'^2 - k^2} dk'^2$$



$$= \tilde{d}(k^2) - ik n(k^2) - \tilde{g} d^R(k^2)$$

$$d_u^R(k^2) = \frac{i}{4k} \left(\log \frac{\mu_+/2 + ik}{\mu_+/2 - ik} - \log \frac{\mu_+/2 + i\eta k}{\mu_+/2 - i\eta k} \right)$$


lhc

$$f^{-1} = \frac{\frac{1}{a} + \frac{1}{2}rk^2}{1 + \tilde{g}(L(k^2) - L_0)} - ik$$

$$d(k^2) = \frac{1}{a} + \frac{1}{2}rk^2 - ik - ik\tilde{g}(L(k^2) - L_0)$$



The left-hand cut



$$t = -2\mathbf{p}^2(1 - \cos \theta)$$

S-wave

$$\frac{1}{2} \int_{-1}^{+1} d\cos \theta \frac{1}{-2\mathbf{p}^2(1 - \cos \theta) - m_\rho^2 + i\varepsilon} = -\frac{1}{4\mathbf{p}^2} \log \left(\frac{4\mathbf{p}^2 + m_\rho^2}{m_\rho^2} + \frac{4\mathbf{p}^2}{m_\rho^4} i\varepsilon \right)$$

