

Nuclear Matter properties from the ladder resummation method

Jose Manuel Alarcón



Universidad
de Alcalá

Works done in collaboration with J. A. Oller

Introduction

Introduction

- There are many different approaches to the calculation of the EoS:

Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory

Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models

Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities

Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities
 - In-medium effects.

Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities
 - In-medium effects
 - Non-standard counting of nucleon propagators.

Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities
 - In-medium effects
 - Non-standard counting of nucleon propagators.
 - Mixing of all partial waves.


Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities
 - In-medium effects
 - Non-standard counting of nucleon propagators.
 - Mixing of all partial waves.

$$\theta(k_F - |\mathbf{k} - \mathbf{a}|)$$

↓

$$\cos \theta \geq \frac{\mathbf{k}^2 + \mathbf{a}^2 - k_F^2}{2|\mathbf{a}||\mathbf{k}|}$$




Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities
 - In-medium effects
 - Non-standard counting of nucleon propagators.
 - Mixing of all partial waves.
 - No regulator.
 - No systematics associated to the choice of a regulator or fine tuning of the scale.

$$\theta(k_F - |\mathbf{k} - \mathbf{a}|)$$

↓

$$\cos \theta \geq \frac{\mathbf{k}^2 + \mathbf{a}^2 - k_F^2}{2|\mathbf{a}||\mathbf{k}|}$$




Introduction

- There are many different approaches to the calculation of the EoS:
 - Ab initio:
 - Quantum Monte Carlo
 - Effective Field Theory
 - Phenomenological:
 - Mean-field models
- We wanted to provide a calculation based on EFT at finite densities
 - In-medium effects
 - Non-standard counting of nucleon propagators.
 - Mixing of all partial waves.
 - No regulator.
 - No systematics associated to the choice of a regulator or fine tuning of the scale.
 - Parameter-free
 - Based on NN observables.

$$\theta(k_F - |\mathbf{k} - \mathbf{a}|)$$

↓

$$\cos \theta \geq \frac{\mathbf{k}^2 + \mathbf{a}^2 - k_F^2}{2|\mathbf{a}||\mathbf{k}|}$$


$\mathbf{a} = \frac{\mathbf{p}_1 + \mathbf{p}_2}{2}$

Formalism

Formalism

- The starting point is the idea of [Kaiser, NPA 860 (2011)] → Resummation of in-medium ladder diagrams to all orders with a contact interaction (scatt. length)

Formalism

- The starting point is the idea of [Kaiser, NPA 860 (2011)] → Resummation of in-medium ladder diagrams to all orders with a contact interaction (scatt. length)
 - The work was extended to include eff. range in S-wave and P-wave → Off-shell effects when reaching the unitary limit.

Formalism

- The starting point is the idea of [Kaiser, NPA 860 (2011)] → Resummation of in-medium ladder diagrams to all orders with a contact interaction (scatt. length)
 - The work was extended to include eff. range in S-wave and P-wave → Off-shell effects when reaching the unitary limit.
- An expansion of the generating functional of in-medium Green function obtained in [Oller, PRC 65 (2002)] provides the diagrams to be resummed in the ladder approximation.

Formalism

- The starting point is the idea of [Kaiser, NPA 860 (2011)] → Resummation of in-medium ladder diagrams to all orders with a contact interaction (scatt. length)
 - The work was extended to include eff. range in S-wave and P-wave → Off-shell effects when reaching the unitary limit.
- An expansion of the generating functional of in-medium Green function obtained in [Oller, PRC 65 (2002)] provides the diagrams to be resummed in the ladder approximation.
- The sum of the Hartree and Fock diagrams gives a contribution due to the interaction of

$$\mathcal{E}_V = \frac{i}{2} \text{Tr} \left(\sum_{n=1}^{\infty} \frac{(t_m L_d)^n}{n} \right) = -\frac{i}{2} \text{Tr} \log [\mathcal{I} - t_m L_d]$$

$$t_m = t_V \sum_{m=0}^{\infty} (L_m t_V)^m = (t_V^{-1} - L_m)^{-1} \quad L_m = -i \int \frac{d^3 k_1}{(2\pi)^3} \theta(k_F - |\vec{k}_1|) \int \frac{d^4 k_2}{(2\pi)^4} \frac{i}{k_2^0 - \frac{|\vec{k}_2|^2}{2m} + i\epsilon} (2\pi)^4 \delta(k_1 + k_2 - 2a) |\vec{k}_1, \vec{k}_2\rangle \langle \vec{k}_1, \vec{k}_2|$$

$$L_d = i \int \frac{d^3 k_1}{(2\pi)^3} \frac{d^3 k_2}{(2\pi)^3} \theta(k_F - |\vec{k}_1|) \theta(k_F - |\vec{k}_2|) |\vec{k}_1 \vec{k}_2\rangle \langle \vec{k}_1 \vec{k}_2|$$

Formalism

- t_m is found to satisfy the following integral equation in partial waves

$$t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\epsilon} V(p', k) [Y^* Y] t_m(k, p)$$

Formalism

- t_m is found to satisfy the following integral equation in partial waves
$$t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\epsilon} V(p', k) [Y^* Y] t_m(k, p)$$
- It can be solved exactly if we consider a contact (momentum dependent interaction)

Formalism

- t_m is found to satisfy the following integral equation in partial waves
$$t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\epsilon} V(p', k) [Y^* Y] t_m(k, p)$$
- It can be solved exactly if we consider a contact (momentum dependent interaction) \longrightarrow Valid up to $k_F \leq \frac{M_\pi}{2}$

Formalism

- t_m is found to satisfy the following integral equation in partial waves
waves
$$t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\varepsilon} V(p', k) [Y^* Y] t_m(k, p)$$
- It can be solved exactly if we consider a contact (momentum dependent interaction) \longrightarrow Valid up to $k_F \leq \frac{M_\pi}{2}$
- Vacuum integrals (no Fermi sea insertions) are regularized through a cutoff (Λ) \longrightarrow Λ is sent to infinity at the end.

Formalism

- t_m is found to satisfy the following integral equation in partial waves
waves
$$t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\epsilon} V(p', k) [Y^* Y] t_m(k, p)$$
- It can be solved exactly if we consider a contact (momentum dependent interaction) \longrightarrow Valid up to $k_F \leq \frac{M_\pi}{2}$
- Vacuum integrals (no Fermi sea insertions) are regularized through a cutoff (Λ) \longrightarrow Λ is sent to infinity at the end.
- We solve the integral equation for t_m with an arbitrary potential (contact terms) $\longrightarrow t_m(\{C_i\}, \Lambda) \longrightarrow t_m(\{a, r, v_2, \dots\}, \Lambda)$

Formalism

- t_m is found to satisfy the following integral equation in partial waves
waves $t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\varepsilon} V(p', k) [Y^* Y] t_m(k, p)$
- It can be solved exactly if we consider a contact (momentum dependent interaction) \longrightarrow Valid up to $k_F \leq \frac{M_\pi}{2}$
- Vacuum integrals (no Fermi sea insertions) are regularized through a cutoff (Λ) \longrightarrow Λ is sent to infinity at the end.
- We solve the integral equation for t_m with an arbitrary potential (contact terms) $\longrightarrow t_m(\{C_i\}, \Lambda) \longrightarrow t_m(\{a, r, v_2, \dots\}, \Lambda)$
- After sending the cutoff to infinity, we recover the t_V on-shell \longrightarrow we can use the experimental data on NN scattering to compute t_m .

$$t_m(\{a, r, v_2, \dots\}) = \left(\tilde{V}^{-1} - L_f - L_m \right)^{-1} = \left(\tilde{t}_V^{-1} - L_m \right)^{-1}$$

Formalism

- t_m is found to satisfy the following integral equation in partial waves
waves $t_m(p', p) = V(p', p) + \frac{m}{(2\pi)^2} \int_0^\infty \frac{k^2 dk}{k^2 - p^2 - i\epsilon} V(p', k) [Y^* Y] t_m(k, p)$
- It can be solved exactly if we consider a contact (momentum dependent interaction) \longrightarrow Valid up to $k_F \leq \frac{M_\pi}{2}$
- Vacuum integrals (no Fermi sea insertions) are regularized through a cutoff (Λ) \longrightarrow Λ is sent to infinity at the end.
- We solve the integral equation for t_m with an arbitrary potential (contact terms) $\longrightarrow t_m(\{C_i\}, \Lambda) \longrightarrow t_m(\{a, r, v_2, \dots\}, \Lambda)$
- After sending the cutoff to infinity, we recover the t_V on-shell \longrightarrow we can use the experimental data on NN scattering to compute t_m .

$$t_m(\{a, r, v_2, \dots\}) = \left(\tilde{V}^{-1} - L_f - L_m \right)^{-1} = \left(\tilde{t}_V^{-1} - L_m \right)^{-1}$$

- Cutoff and regulator-independent results.

Results

Results



- Unitary limit [J. M. Alarcón, J. A. Oller, AOP 437 (2022)]

$$\mathcal{E}_V = \frac{8k_F^5}{m\pi^3} \int_0^1 s^2 ds \int_0^{\sqrt{1-s^2}} \kappa d\kappa \arctan \left(\frac{a_0 k_F I(s, \kappa)}{1 - a_0 k_F R(s, \kappa)/\pi} \right)$$

$$\mathcal{E}_V = \frac{8k_F^5}{m\pi^3} \int_0^1 ds s^2 \int_0^{\sqrt{1-s^2}} d\kappa \kappa \arctan \left(\frac{a_0 k_F I(s, \kappa)}{1 - a_0 r_0 k_F^2 \kappa^2 / 2 - a_0 k_F R(s, \kappa)/\pi} \right)$$

$$\xi = 1 - \frac{80}{\pi} \int_0^1 ds s^2 \int_0^{\sqrt{1-s^2}} d\kappa \kappa \arctan \left(\frac{\pi I(s, \kappa)}{R(s, \kappa)} \right) = 0.5066$$


$$\xi(k_F) = 1 - \frac{80}{\pi} \int_0^1 ds s^2 \int_0^{\sqrt{1-s^2}} d\kappa \kappa \arctan \left(\frac{\pi I(s, \kappa)}{\pi r_0 k_F^2 \kappa^2 / 2 + R(s, \kappa)} \right)$$



Results



- Cold atoms [J. M. Alarcón, J. A. Oller, PRC 106 (2022)]

Spin-balanced fermionic quantum liquid with P-wave interactions



Results

- Symmetric Nuclear Matter [J. M. Alarcón, J. A. Oller, PRC 107 (2023)]




- To estimate the uncertainty:

- Multiply the off-shell dependence of Lm times the gaussian regulator

$$\exp\left[-\frac{(k - M_\pi/2)^2}{\lambda^2}\right] \quad \lambda > M_\pi$$

- Consider a density-dependence nucleon mass



$$m_N(n) = \frac{m_N}{1 + \frac{n}{n_s} \left(\frac{m_N}{m_N^*} - 1\right)} \quad m_N^* \in [0.7m_N, 0.9m_N]$$



Results

- Pure Neutron Matter

[J. M. Alarcón, J. A. Oller, PRC 107 (2023)]




- To estimate the uncertainty:

- Multiply the off-shell dependence of L_m times the gaussian regulator

$$\exp\left[-\frac{(k - M_\pi/2)^2}{\lambda^2}\right] \quad \lambda > M_\pi$$

- Consider a density-dependence nucleon mass

$$m_N(n) = \frac{m_N}{1 + \frac{n}{n_s} \left(\frac{m_N}{m_N^*} - 1\right)} \quad m_N^* \in [0.7m_N, 0.9m_N]$$



Results

- To study the properties of nuclear matter at saturation density we use the parametrization [Gandolfi et al., Mon. Not. Roy. Astron. Soc. 404 (2010)]

$$S(n) = (E/A)_{PNM}(n) - (E/A)_{SNM}(n) \quad S(n) = C_s \left(\frac{n}{n_s} \right)^{\gamma_s} \quad S_0 = S(n_s) \quad L = 3n_s \left. \frac{dS(n)}{dn} \right|_{n_s}$$

Results

- To study the properties of nuclear matter at saturation density we use the parametrization [Gandolfi et al., Mon. Not. Roy. Astron. Soc. 404 (2010)]

$$S(n) = (E/A)_{PNM}(n) - (E/A)_{SNM}(n) \quad S(n) = C_s \left(\frac{n}{n_s} \right)^{\gamma_s} \quad S_0 = S(n_s) \quad L = 3n_s \left. \frac{dS(n)}{dn} \right|_{n_s}$$


- We fit this parametrization to our results at low densities.

$$C_s = 34.77(15) \text{ MeV}$$

$$31.10 \leq S_0 \leq 36.57 \text{ MeV}$$


$$\gamma_s = 0.667(3)$$

$$57.82 \leq L \leq 78.29 \text{ MeV}$$



Results



- Using astrophysical data to constrain the EoS



[J. M. Alarcón, E. Lope-Oter and J. A. Oller, arXiv: 2410.14776]

Results




- Using astrophysical data to constrain the EoS



[J. M. Alarcón, E. Lope-Oter and J. A. Oller, arXiv: 2410.14776]

Results




- Using astrophysical data to constrain the EoS




[J. M. Alarcón, E. Lope-Oter and J. A. Oller, arXiv: 2410.14776]

Results

- Using astrophysical data to constrain the EoS



$$M_{\text{TOV}} \leq 2.7 M_{\odot}$$



[J. M. Alarcón, E. Lope-Oter and J. A. Oller, arXiv: 2410.14776]

Future applications


- A reliable EoS is important for multiple applications.

Future applications

- A reliable EoS is important for multiple applications.
- Dark Matter searches in Neutron Stars


Future applications

- A reliable EoS is important for multiple applications.
- Dark Matter searches in Neutron Stars



Future applications


- A reliable EoS is important for multiple applications.
- Dark Matter searches in Neutron Stars



HESS J1731-347
(Possible admixture
of NS+DM)

Future applications

- A reliable EoS is important for multiple applications.
- Dark Matter searches in Neutron Stars




Different DM fractions

HESS J1731-347
(Possible admixture
of NS+DM)

Summary and Conclusions

Summary and Conclusions

- We develop a method to compute the energy per particle in nuclear matter at low densities.
- E/A is directly related to in-vacuum NN scattering amplitude.
- Regularizator-independent results
- Can be used to study:
 - Unitary limit
 - Cold atoms
 - Equation of state of symmetric nuclear matter and pure neutron matter.
- EoS of PNM constraints equation of state of neutron stars at low densities  Determination of the EoS of a NS
 - Astrophysical applications.
 - Dark Matter searches in compact objects.

FIN