


Lattice QCD studies on meson-nucleon interactions and genuine pentaquarks

Kotaro Murakami (村上 耕太郎)

Institute of Science Tokyo/RIKEN iTHEMS



based on

KM, S. Aoki, T. Doi, Y. Lyu, W. Yamada (HAL QCD), accepted in PRD [arXiv: 2509.00838]



W. Yamada, Y. Lyu, KM, T. Doi, (HAL QCD), in preparation

核力・ハドロン間相互作用と量子多体計算の進展, 京都大学, 2026年2月16-18日



Hadron interactions from lattice QCD

- our goal: understand exotic hadrons/nuclei/matters via hadron interactions using lattice QCD


Lattice QCD



hadron interactions (scatterings)



Exotic hadrons



(Exotic) Nuclei



Nuclear matter, Neutron stars

- Finite-volume method [Lüscher, 1991]
- HAL QCD method [Ishii, Aoki, Hatsuda 2007]

HAL QCD method

- extract an **interaction potential**

from Nambu-Bethe-Salpeter (NBS) wave functions:

$$\Psi^W(\mathbf{r}) = \langle 0 | \hat{O}_1(\mathbf{r}) \hat{O}_2(\mathbf{0}) | 1, 2; W \rangle \xrightarrow{r \rightarrow \infty} A \frac{\sin(kr - \frac{l}{2}\pi + \delta^l(k))}{kr} e^{i\delta^l(k)} Y_{l,m}(\Omega)$$

$$\longrightarrow \int d^3r' \boxed{U(\mathbf{r}, \mathbf{r}')} \Psi^W(\mathbf{r}') = \left(\frac{k^2}{2\mu} + \frac{\nabla^2}{2\mu} \right) \Psi^W(\mathbf{r})$$

interaction potential

[Ishii, Aoki, Hatsuda, PRL (2007)]
[Aoki, Hatsuda, Ishii, PTP (2010)]

- practical calc. : **time-dependent HAL QCD method** [Ishii et al., PLB (2012)]

+ **local approximation** ($U(\mathbf{r}, \mathbf{r}') \approx V(\mathbf{r}) \delta^{(3)}(\mathbf{r} - \mathbf{r}')$)

$$R(\mathbf{r}, t) = \frac{\langle O_1(\mathbf{r}, t) O_2(\mathbf{0}, t) \bar{J}(0) \rangle}{\langle O_1(t) \bar{O}_1(0) \rangle \langle O_2(t) \bar{O}_2(0) \rangle} \approx \sum_n C_{\bar{J}, n} \underline{\Psi^{W_n}(\mathbf{r})} e^{-(W_n - m_1 + m_2)t}$$

NBS wave function

$$\longrightarrow V(\mathbf{r}) \simeq \frac{1}{R(\mathbf{r}, t)} \left(\frac{\nabla^2}{2\mu} - \frac{\partial}{\partial t} + \frac{1}{8\mu} \frac{\partial^2}{\partial t^2} \right) R(\mathbf{r}, t)$$

Current situation of HAL QCD studies

- ~2024: near-physical-point simulations ($m_\pi \approx 146$ MeV)

- $\Omega\Omega$ [Gongyo et al., 2018]
- $N\Omega$ [Iritani et al., 2019]
- nucleon/hyperon interactions: [S. Aoki and T. Doi, Front. in Phys. **8**, 307 (2020) (review)]
- H dibaryon [Sasaki et al., 2020]
- $\Omega_{ccc}\Omega_{ccc}$ [Lyu et al., 2021]
- T_{cc}^+ [Lyu et al., 2023]
- $N\phi, NJ/\psi$ [Lyu et al., 2022]
[Lyu et al., 2025]

- **2025~: on-physical-point simulations** ($m_\pi \approx 137$ MeV)

- configuration generation: done
[HAL QCD Collab., 2024]
- $N\Omega_{ccc}$ [Zhang et al., 2025]
- KN [KM, Aoki, Doi, Lyu, Yamada, [arXiv: 2509.00838]]
- $\bar{D}N$ [Yamada, Lyu, KM, Doi, in preparation]
- YN (Talk by Murase-san today)
- $\Lambda\Lambda, \Xi N$ [Sasaki et al., in preparation]
- $N\Omega$ [Zhang et al., work in progress]

Current situation of HAL QCD studies

- ~2024: near-physical-point simulations ($m_\pi \approx 146$ MeV)

- $\Omega\Omega$ [Gongyo et al., 2018]
- $N\Omega$ [Iritani et al., 2019]
- nucleon/hyperon interactions: [S. Aoki and T. Doi, Front. in Phys. **8**, 307 (2020) (review)]
- H dibaryon [Sasaki et al., 2020]
- $\Omega_{ccc}\Omega_{ccc}$ [Lyu et al., 2021]
- T_{cc}^+ [Lyu et al., 2023]
- $N\phi, NJ/\psi$ [Lyu et al., 2022]
[Lyu et al., 2025]

- **2025~: on-physical-point simulations** ($m_\pi \approx 137$ MeV)

- configuration generation: done [HAL QCD Collab., 2024]

- $N\Omega_{ccc}$ [Zhang et al., 2025]

- KN [KM, Aoki, Doi, Lyu, Yamada, [arXiv: 2509.00838]]

- $\bar{D}N$ [Yamada, Lyu, KM, Doi, in preparation]

- YN (Talk by Murase-san today)

- $\Lambda\Lambda, \Xi N$ [Sasaki et al., in preparation]

- $N\Omega$ [Zhang et al., work in progress]

- other on-going HAL QCD projects

- hyperon int. from HAL QCD

(Talk by Inoue-san today)

- negative-parity nuclear force

(Talk by Sugiura-san yesterday)

- optimized operator constructions

[Yan et al., [arXiv:2507.09933]]

[Yan et al., [arXiv:2507.09930]]

- $\Lambda(1405)$ in SU(3) limit

[KM, Aoki, in preparation]

Contents


- Introduction
- Motivation for $KN/\bar{D}N$ interactions
- Numerical results
 - KN interactions
KM, S. Aoki, T. Doi, Y. Lyu, W. Yamada (HAL QCD), [arXiv: 2509.00838]
 - $\bar{D}N$ interactions
W. Yamada, Y. Lyu, KM, T. Doi (HAL QCD), in preparation

Contents

- Introduction
- Motivation for $KN/\bar{D}N$ interactions
- Numerical results
 - KN interactions
KM, S. Aoki, T. Doi, Y. Lyu, W. Yamada (HAL QCD), [arXiv: 2509.00838]
 - $\bar{D}N$ interactions
W. Yamada, Y. Lyu, KM, T. Doi (HAL QCD), in preparation

Motivation for meson-nucleon interactions (1/2)

- interactions between a nucleon and $q\bar{Q}$ -type meson



- no quark-antiquark pair creation/annihilation


- can provide **understandings of quark/hadron dynamics**

- restoration of chiral symmetry in medium
- possible existence of genuine pentaquark

1. restoration of chiral symmetry in medium

- $\langle \bar{u}u + \bar{s}s \rangle^*(\rho_N)$ from KN scattering data
[Iizawa, Jido, Hübsch, PTEP (2024)]
- rich scattering experimental data since 1960s
- no data near threshold** due to the experimental difficulty


➔ **large ambiguity** (especially for $l=0$)




[Iizawa, Jido, Hübsch, PTEP (2024)]

Motivation for meson-nucleon interactions (2/2)

2. possible existence of open-strange/charm genuine pentaquark




- Θ^+ ($uudd\bar{s}$) pentaquark in KN
 - most experiments after the LEPs gave negative results [LEPS, PRL (2003)]
 - **no definitive conclusion from theoretical side**



[LEPS, PRL (2003)]

- Θ_c^+ ($uudd\bar{c}$) pentaquark in $\bar{D}N$
 - $\bar{D}N$ would be attractive due to $\bar{D}^*N-\bar{D}N$ coupling
 → bound state?





- scarce experimental data for $\bar{D}N$ (only by ALICE) [ALICE, PRD (2022)]
- **phenomenological-model prediction is uncertain**

Model	$\bar{D}N(I = 0)$	$\bar{D}N(I = 1)$
SU(4) contact [185]	None	None
SU(8) contact [193]	2805	None
Meson exchange [194]	None	None
Pion exchange [192]	2804	None
Chiral quark model [195]	None	None

[Hosaka et al., Prog. Part. Nucl. Phys. (2017)]

$KN/\bar{D}N$ interactions in lattice QCD



- $KN/\bar{D}N$: relatively easy to calculate
 - no annihilation diagram \rightarrow cheap computational cost
 - no need for coupled-channel analysis
 - **physical point simulations are now possible** (for S-wave component)
($m_\pi \approx 137$ MeV) [HAL QCD, PRD (2024)]

Lattice QCD compensates the missing profiles of $KN/\bar{D}N$

- this talk: theoretical study on $KN/\bar{D}N$ interactions using **lattice QCD on physical point**
 - how does the $KN/\bar{D}N$ **interactions** behave?
 - what are $KN/\bar{D}N$ scatterings like **near the threshold**?
 - does the **genuine pentaquarks** exist?

Calculation strategy

- u, d, s quarks: $N_f = 2 + 1$ gauge configurations ($m_u = m_d < m_s$)

[HAL QCD, PRD (2024)]


on the physical point

- clover fermion + Iwasaki gauge w/ stout smearing
- $V = (96a)^4 \approx (8.1 \text{ fm})^4$, $a = 0.084372(54)({}_{-6}^{+109}) \text{ fm}$
- $m_\pi \approx 137 \text{ MeV}$, $m_K \approx 502 \text{ MeV}$, $m_N \approx 940 \text{ MeV}$
- c quark: quench approximation for the sea quark, action w/ physical mass for the valence quark
- (approximated) S-wave component of wave functions are extracted


Contents

- Introduction
- Motivation for $KN/\bar{D}N$ interactions
- Numerical results
 - KN interactions
KM, S. Aoki, T. Doi, Y. Lyu, W. Yamada (HAL QCD), [arXiv: 2509.00838]
 - $\bar{D}N$ interactions
W. Yamada, Y. Lyu, KM, T. Doi (HAL QCD), in preparation


KN local potentials



$l = 1$






$l = 0$



- $l=1$: repulsive core
- $l=0$: repulsive core + attractive pocket

KN Phase shifts





- black plots: experimental results
- gray lines: partial-wave analysis (parameterization method)

• no bound states or resonances found

• larger in magnitude than experimental data for $I=1$


- more careful determination is required both in experiment and theory
- necessary to see the lattice discretization effect in future

S-wave total cross sections



- color bands: our results (**only S-wave component**)
- black plots: experimental results (**including all partial waves**)

- $I=0$: consistent with (u)ChPT [Aoki, Jido, PTEP (2019)]
 → $I=0$ cross section is dominant by P-wave?



Contents

- Introduction

- Motivation for $KN/\bar{D}N$ interactions

- Numerical results




 - KN interactions

KM, S. Aoki, T. Doi, Y. Lyu, W. Yamada (HAL QCD), [arXiv: 2509.00838]

 - $\bar{D}N$ interactions




W. Yamada, Y. Lyu, KM, T. Doi (HAL QCD), in preparation

$\bar{D}N$ local potentials




- repulsive core + attractive pocket for both isospins




Comparison between $\bar{D}N$ and KN



- $\bar{D}N$ is more attractive than KN
- sign of attraction attributed to coupling of \bar{D}^*N ?




$\bar{D}N$ Phase shifts




- no bound states or resonances found
- more attractive behavior compared with KN

Comparison with $\bar{D}N$ phenomenological studies



- scattering lengths a_0

$$k \cot \delta_0(k) = \frac{1}{a_0} + \frac{1}{2} r_{\text{eff}} k^2 + \mathcal{O}(k^4)$$



- same order as those in the models that do not exhibit bound state
- our results can constrain the model parameters?

Summary

- we study S-wave $KN/\bar{D}N$ interactions in lattice QCD using HAL QCD method on the physical point
- KN interactions exhibit repulsive behavior in short distance
- KN $I = 1$ phase shifts are larger than experiment
- $\bar{D}N$ is more attractive than KN , which may be a sign of attraction attributed to coupling of \bar{D}^*N
- our lattice results
- **no resonance or bound state in KN and $\bar{D}N$**

Future prospects

- (KN) need estimation of systematics by simulation w/ different lattice spacing
- $(\bar{D}N)$ \bar{D}^*N - $\bar{D}N$ coupled channel analysis might change the results
- $(KN, \bar{D}N)$ P-wave calculations (at heavier quark masses)
- meson-nucleon w/ a bottom quark ($BN - B^*N$)