

t-channel 特異点に対する 解析的処方

横浜国立大学 博士課程 1 年、理研仁科センター

佐藤龍政

This talk is based on arXiv:2505.10890.

共同研究者：浅井健人(京大基研)、
広島渚(横浜国大、理研iTHEMS)、
佐藤丈(横浜国大)、梁正樹(埼玉大)

素粒子物理学の進展2025

京都大学基礎物理学研究所 2025/9/1~9/5

発表日(2025/9/1)

Plan

- Introduction
- Analytic prescription
- Application : $L_\mu - L_\tau$ model
majoron production in the early universe
- Summary

Introduction

Interaction in QFT

model building (construct Lagrangian)




Check the Lagrangian by scattering



scattering amplitude $|\mathcal{M}|^2$ is important

Divergence in scattering amplitude



propagator $\frac{1}{p^2 - m^2 + i\varepsilon}$

- Off-shell (Virtual)

$$p^2 - m^2 \neq 0 \rightarrow \text{finite}$$


- On-shell

$$p^2 - m^2 = 0 \rightarrow \text{divergence}$$

$$\sim \frac{1}{i\varepsilon}$$

Physical interpretation


s-channel



Pole of s-channel = inverse decay $\times$ decay

➔ Mediator is **unstable**


t-channel



Pole of t-channel = decay $\times$ inverse decay

➔ Mediator **can** be **stable**

s-channel amplitude




Inverse decay and decay
→ Intermediate particles
are unstable

Tree propagator

$M \text{ ————— } M^*$

$$\frac{1}{p^2 - m^2 + i\varepsilon}$$

One-loop propagator




$$\frac{1}{p^2 - m^2 + im\Gamma}$$

Imaginary part is decay width Γ **Breit-Wigner distribution**
 propagator is modified by finite decay width!

Does not diverge at the pole.

Resonance rather than singularity

t-channel amplitude



Decay and Inverse decay
 → Intermediate particles
 can be stable.

Tree propagator

$$M \text{ ————— } M^*$$

$$\frac{1}{p^2 - m^2 + i\varepsilon}$$

Stable mediator → no decay width!

true singularity ?

How can we regularize the amplitude ?

Previous works & our focus

Previous work

- introducing effective scattering width (collider)
K. Melnikov, G. L. Kotkin, and V. G. Serbo. PRD, 54:3289–3295, 1996.
- Just to ignore as it(divergent part) is included in inverse decay(cosmology)
M. Escudero, Dan Hooper, G. Krnjaic, M. Pierre. JHEP, 03:071, 2019
- numerical subtraction of divergent part
Kento Asai, Tomoya Asano, Joe Sato, and Masaki J. S. Yang. PTEP, 2024(7):073E01, 2024.

Previous works & our focus

Previous work

- Not applicable to cosmology

K. Melnikov, G. L. Kotkin, and V. G. Serbo. PRD, 54:3289–3295, 1996.

- Is it realistic?

M. Escudero, Dan Hooper, G. Krnjaic, M. Pierre. JHEP, 03:071, 2019



- High computational cost

Kento Asai, Tomoya Asano, Joe Sato, and Masaki J. S. Yang. PTEP, 2024(7):073E01, 2024.

Analytic prescription

Structure of divergence

$$\sigma = \sigma_{\text{off-shell}} + \sigma_{\text{on-shell}}$$



$$\sigma_{\text{on-shell}} = \frac{1}{\varepsilon} E \cdot \Gamma(\phi_3 \rightarrow \Phi \phi_1) \cdot \sigma(\phi_4 \Phi \rightarrow \phi_2)$$

The on-shell contribution is a divergence of $1/\varepsilon$.

Structure of divergence

Calculation of t-channel singularity part

$$\begin{aligned}
 \int_a^b dX |\mathcal{M}^t|^2 &\sim \int_a^b dX \frac{1}{X^2 + \epsilon^2} & X &= (p_3 - p_2)^2 - m^2 \\
 \int_{a/\epsilon}^{b/\epsilon} d(\epsilon \bar{X}) \frac{1}{\epsilon^2} \frac{1}{\bar{X}^2 + 1} &= \frac{1}{\epsilon} \left(\arctan\left(\frac{b}{\epsilon}\right) - \arctan\left(\frac{a}{\epsilon}\right) \right) \\
 &= \frac{1}{a} - \frac{1}{b} + \mathcal{O}(\epsilon) & \text{If } ab > 0 & \text{No on-shell terms} \\
 &= -\boxed{\frac{\pi}{\epsilon}} + \frac{1}{a} - \frac{1}{b} + \mathcal{O}(\epsilon) & \text{If } a < 0 < b & \text{on-shell terms}
 \end{aligned}$$

on-shell term that should be subtracted

To avoid double counting : already included
by **Decay & Inverse Decay**

Application

Majoron production ($L_\mu - L_\tau$ model)

- Interaction of Z'

e.g. Phys.Rev.D 100 (2019) 9, 095012

$$\mathcal{L}_{Z'} = -\frac{1}{4}Z'^{\rho\sigma}Z'_{\rho\sigma} + \frac{1}{2}m_Z^2 Z'^\rho Z'_\rho + g_{\mu-\tau} Z'_\rho J_{\mu-\tau}^\rho$$

$$Z'_{\rho\sigma} \equiv \partial_\rho Z'_\sigma - \partial_\sigma Z'_\rho$$

$$J_{\mu-\tau}^\rho \equiv \bar{\mu}\gamma^\rho\mu + \bar{\nu}_\mu\gamma^\rho P_L \nu_\mu - \bar{\tau}\gamma^\rho\tau - \bar{\nu}_\tau\gamma^\rho P_L \nu_\tau$$

- Interaction of majoron ϕ (pNGB)

$$\mathcal{L}_\phi = h_{\alpha\beta} \bar{\nu}_{L,\alpha} \nu_{L,\beta}^c \phi + \text{H.c.}$$

decay & inverse decay

$$\begin{aligned} Z' &\leftrightarrow \nu_{\alpha'} \bar{\nu}_{\alpha'} \quad (\alpha' = \mu, \tau) \\ \phi &\leftrightarrow \nu_\alpha \nu_\beta \end{aligned}$$

scattering


$$\begin{aligned} Z' \nu_\alpha &\leftrightarrow \phi \bar{\nu}_\beta \quad (\nu \leftrightarrow \bar{\nu}) \\ Z' \phi &\leftrightarrow \nu_\alpha \nu_\beta \quad (\nu \leftrightarrow \bar{\nu}) \end{aligned}$$

Majoron production in the early universe


Majoron (pNGB of lepton number) may have been created in the early universe.
→ The effective d.o.f. of neutrinos and cosmic expansion rate may change.

Let's discuss $U(1)_{L_\mu - L_\tau}$ model

(inverse) decay



$Z' - \phi$ scattering (New!)



M. Escudero, Dan Hooper, G. Krnjaic, M. Pierre.
JHEP, 03:071, 2019

t-channel singularity in Majoron production


$Z' \nu_\alpha \leftrightarrow \phi \bar{\nu}_\beta$ in t-channel diagram

Intermediate particle can be on-shell

→ Amplitude. $|\mathcal{M}|^2$ is **divergent**

$$|\mathcal{M}_{Z' \nu \leftrightarrow \phi \bar{\nu}}^t|^2 \propto \frac{1}{(p_3 - p_2)^4 + \varepsilon^2}$$

Regularized by our new method



Net production rate

Generation of Majoron ϕ in the early universe is evaluated by Boltzmann eq.

$$\frac{dn_a}{dt} = -3Hn_a + \frac{\delta n_a}{\delta t}$$

thermal averaged event rates of scattering and inverse decay

$$\langle \Gamma_S \rangle = \frac{1}{n_{\phi,eq}} \left. \frac{\delta n_{\phi}}{\delta t} \right|_S, \quad \langle \Gamma_{ID} \rangle = \frac{1}{n_{\phi,eq}} \left. \frac{\delta n_{\phi}}{\delta t} \right|_{ID}, \quad d\Pi_a = \frac{d^3 \mathbf{p}_a}{(2\pi)^3 2E_a}$$

$$\left. \frac{\delta n_a}{\delta t} \right|_{ID} = -\frac{1}{S_{ij}} \int d\Pi_a \int d\Pi_i d\Pi_j (2\pi)^4 \delta^{(4)}(p_a - p_i - p_j) \sum_{\text{spins}} |\mathcal{M}_{a \leftrightarrow ij}|^2 \Lambda_{ID} \{f_i^{\text{MB}}\}$$

$$\left. \frac{\delta n_a}{\delta t} \right|_S = - \int d\Pi_a d\Pi_b d\Pi_c d\Pi_d \Lambda_S \{f_i^{\text{MB}}\} (2\pi)^4 \delta^{(4)}(\mathbf{p}_a + \mathbf{p}_b - \mathbf{p}_c + \mathbf{p}_d) \sum_{\text{spins}} |\mathcal{M}_{ab \leftrightarrow cd}|^2$$

Calculation procedure

$$\left. \frac{\delta n_a}{\delta t} \right|_S = - \int d\Pi_a d\Pi_b d\Pi_c d\Pi_d \Lambda_S \{f_i^{\text{MB}}\} (2\pi)^4 \delta^{(4)}(\mathbf{p}_a + \mathbf{p}_b - \mathbf{p}_c + \mathbf{p}_d) \sum_{\text{spins}} |\mathcal{M}_{ab \leftrightarrow cd}|^2$$

$$I = \int d\Pi_3 d\Pi_4 (2\pi)^4 \delta^{(4)}(p_1 + p_2 - p_3 - p_4) \sum_{\text{spins}} |\mathcal{M}_{\nu_\alpha \phi \leftrightarrow Z' \nu_\beta}|^2$$

$$I^t = \frac{16g_\beta^2 |h_{\alpha\beta}|^2}{2\pi\sqrt{s}p_2} \theta(\sqrt{s} - m_{Z'}) \left[\frac{1}{\varepsilon} \tan^{-1} \frac{x}{\varepsilon} m_{Z'}^2 (s - 2m_{Z'}^2 + 2p_2\sqrt{s}) \right] d\Pi_a = \frac{d^3\mathbf{p}_a}{(2\pi)^3 2E_a}$$


By on-shell subtraction

$$- \frac{1}{2} \log(x^2 + \varepsilon^2)(s - 2m_{Z'}^2) + \left(x - \varepsilon \tan^{-1} \frac{x}{\varepsilon} \right) \frac{1}{2m_{Z'}^2} (m_{Z'}^2 - 2p_2\sqrt{s} - x) \Bigg]$$

$$I_0 = \frac{16g_\beta^2 |h_{\alpha\beta}|^2}{2\pi\sqrt{s}p_2} \theta(\sqrt{s} - m_{Z'}) \left[-\frac{1}{x} m_{Z'}^2 (s - 2m_{Z'}^2 + 2p_2\sqrt{s}) \right.$$

$$\left. - \frac{1}{2} \log(x^2 + \varepsilon^2)(s - 2m_{Z'}^2) + \frac{x}{2m_{Z'}^2} (m_{Z'}^2 - 2p_2\sqrt{s} - x) \right]$$

Production efficiency



$$\langle \Gamma_S \rangle = \frac{1}{n_{\phi, \text{eq}}} \left. \frac{\delta n_\phi}{\delta t} \right|_S, \quad \langle \Gamma_{\text{ID}} \rangle = \frac{1}{n_{\phi, \text{eq}}} \left. \frac{\delta n_\phi}{\delta t} \right|_{\text{ID}},$$

Gamov Criterion

$$\langle \Gamma_{S, \text{ID}} \rangle / H > 1$$

For Majoron
to be thermalized !

Majoron ϕ can have been made a lot in the early universe?
→ previous works could underestimate Majoron effects



Summary

- Generally, divergence occurs when intermediate particles become on-shell.
 - s-channel: regulated by the decay width.
 - t-channel: divergence can appear.
- In this study, we proposed an analytical method to eliminate the t-channel singularity.
 - divergent part $\sim 1/\varepsilon$: subtracted
real particle contribution = decay & Inverse decay
 - scattering part : remains

Future Work

- We plan to perform parameter scan for cosmological systems.
 - Constraints on new particles such as Majoron can be more stringent.


- We apply our method to $\mu^+ \mu^- \rightarrow \bar{\nu} e^- W^+$



Back up

Structure of divergence

$$\begin{aligned}
 \sigma &= \frac{y_1^2 y_2^2}{2E_3 2E_4} \int d\Pi_1 d\Pi_2 (2\pi)^4 d^4 p \delta^{(4)}(p_3 - p - p_2) \delta^{(4)}(p + p_4 - p_1) \left| \frac{i}{p^2 - m^2 + i\varepsilon} \right|^2 \\
 &= \frac{y_1^2 y_2^2}{2E_3 2E_4} \int d\Pi_1 d\Pi_2 (2\pi)^4 \text{p.v.} \Big|_{p^2 \neq m^2} \left(\int d^4 p \delta^{(4)}(p_3 - p - p_2) \delta^{(4)}(p + p_4 - p_1) \frac{1}{(p^2 - m^2)^2} \right) \\
 &\quad + \frac{y_1^2 y_2^2}{2E_3 2E_4} \lim_{\eta \rightarrow 0^+} \int d\Pi_1 d\Pi_2 (2\pi)^4 \int_{m^2 - \eta < p^2 < m^2 + \eta} d^4 p \delta^{(4)}(p_3 - p - p_2) \delta^{(4)}(p + p_4 - p_1) \frac{1}{(p^2 - m^2)^2 + \varepsilon^2} \\
 &\equiv \sigma_{\text{off-shell}} + \sigma_{\text{on-shell}} ,
 \end{aligned}$$



$$\begin{aligned}
 \sigma_{\text{on-shell}} &= \frac{y_1^2 y_2^2}{2E_3 2E_4} \int d\Pi_1 d\Pi_2 (2\pi)^4 \int_{m^2 - \eta < p^2 < m^2 + \eta} d^4 p \delta^{(4)}(p_3 - p - p_2) \delta^{(4)}(p + p_4 - p_1) \frac{\pi}{\varepsilon} \frac{1}{\pi} \frac{\varepsilon}{(p^2 - m^2)^2 + \varepsilon^2} \\
 &= \frac{y_1^2 y_2^2}{2E_3 2E_4} \int d\Pi_1 d\Pi_2 (2\pi)^4 \int_{m^2 - \eta < p^2 < m^2 + \eta} d^4 p \delta^{(4)}(p_3 - p - p_2) \delta^{(4)}(p + p_4 - p_1) \frac{\pi}{\varepsilon} \delta(p^2 - m^2) \\
 &= \frac{\pi}{\varepsilon} \frac{y_1^2 y_2^2}{2E_3 2E_4} \int d\Pi_1 d\Pi_2 \frac{d^3 \mathbf{p}}{2E} (2\pi)^4 \delta^{(4)}(p_3 - p - p_2) \delta^{(4)}(p + p_4 - p_1) \\
 &= \frac{1}{\varepsilon} E \cdot \frac{y_1^2}{2E_3} \int d\Pi_2 \frac{d^3 \mathbf{p}}{2E (2\pi)^3} (2\pi)^4 \delta^{(4)}(p_3 - p - p_2) \cdot \frac{y_2^2}{2E_4 2E} \int d\Pi_1 (2\pi)^4 \delta^{(4)}(p + p_4 - p_1) \\
 &= \frac{1}{\varepsilon} E \cdot \Gamma(\phi_3 \rightarrow \Phi \phi_1) \cdot \sigma(\phi_4 \Phi \rightarrow \phi_2) .
 \end{aligned}$$

