



MINISTERIO
DE CIENCIA, INNOVACIÓN
Y UNIVERSIDADES



Cofinanciado por
la Unión Europea



AGENCIA
ESTATAL DE
INVESTIGACIÓN



Institut de Ciències del Cosmos
UNIVERSITAT DE BARCELONA



EXCELENCIA
MARÍA
DE MAEZTU
04/2025-03/2031

The neutron skin thickness in atomic nuclei

Xavier Roca-Maza

Intersection of nuclear structure and
high-energy nuclear collisions 2026

April 13th–24th de 2026
Yukawa Institute for Theoretical Physics




UNIVERSITAT DE
BARCELONA



Neutron skin thickness in atomic nuclei

What is it?



The **mean square radius** of the **neutron** (n) or **proton** (p) **density distribution** ($\rho_q(\vec{r})$ with $q = n, p$) is defined as:


$$\langle r_q^2 \rangle = \int d\vec{r} r^2 \rho_q(\vec{r})$$

The neutron skin thickness is defined as the difference between the neutron and proton radii

$$\Delta r_{np} \equiv \langle r_n^2 \rangle^{1/2} - \langle r_p^2 \rangle^{1/2}$$


For **atomic nuclei** with $N > Z$ we expect a **neutron skin**, **larger** as **larger** the average repulsion (**pressure**) felt by the neutrons

In **first approximation** (within a Local Density Approximation) **the pressure** that a **neutron uniform system feels** at the densities typical for the atomic nucleus



Neutron skin thickness in atomic nuclei

An important physical insight!




Credit: FrankRamsrott

Neutron skin thickness is a proxy to the pressure (nuclear force) of an ideal uniform system composed only by neutrons and protons (L)


Droplet Model:

$$\Delta r_{np} \approx \frac{1}{12} \frac{N - Z}{A} \frac{R}{J} L$$

State-of-the-art microscopic models:



$$\Delta \tilde{r}_{np} \equiv \langle r_n^2 \rangle^{1/2} - \langle r_p^2 \rangle^{1/2}$$



Model predictions are very spread

Neutron skin thickness in atomic nuclei

Why is it important?

Constraints the nuclear equation of state (EoS)

The neutron skin is strongly linked to the density dependence of the energy per particle in a uniform system

[WE WILL DISCUSS THIS IN WHAT FOLLOWS]

Connects nuclei to neutron stars

The same physics governing neutron skins also determines properties of neutron stars related to their size and deformability (quadruple polarizability) as well as how exotic are nuclei present in their crusts.

Tests nuclear structure models

Precise measurements discriminate between competing theoretical models, improving our understanding of nuclear forces.

Impacts reactions and stability of exotic nuclei

Neutron distribution affects reaction cross sections, decay properties, and predictions for very neutron-rich nuclei (relevant for rare-isotope physics).

Informs r-process nucleosynthesis

Better constraints on neutron-rich matter improve models of heavy element formation in astrophysical environments.

Provides electroweak physics tests

Techniques like parity-violating electron scattering probe neutron distributions with minimal strong-interaction uncertainties, offering clean tests of the Standard Model in nuclei.

Nuclear Equation of State (EoS)

What is it? How is defined?

Definition: the energy (E) per nucleon ($A=N+Z$), $e \equiv E/A$, of an **uniform system of neutrons (N) and protons (Z)** as a function of the **neutron ($\rho_n = N/V$) and proton ($\rho_p = Z/V$) densities ...**

→ ... **at zero temperature:** room temperature $10^2\text{K} \rightarrow 10^{-8}$ MeV while “cold” neutron stars are at about $10^{10}\text{K} \rightarrow 1$ MeV. **Separation energy** in stable nuclei (equivalent to ionization energy in atoms) is of **several MeV**.

→ ... **unpolarized:** energy favors **couples** of neutrons and protons **occupying the same state but with opposite spins** (equivalent to electrons in atoms)

→ ... **isospin symmetric:** neutron-neutron, proton-proton and neutron-proton **nuclear interaction** are very **similar** among them. **Masses** of neutrons and protons are almost **degenerate**. Hence neutrons and protons can be thought as **two states** of the **same particle** with different isobaric spin or **isospin** (in analogy with spin): the **nucleon**.

→ ... and **no Coulomb:** **idealized uniform system** (focus on strong interaction, much stronger than Coulomb at nuclear scale $\sim 10^{15}$ fm). Real systems are finite and electrically neutral so no divergences in adding Coulomb.

Nuclear Equation of State (EoS)

What is it? How is defined?

Definition: the energy (E) per nucleon ($A=N+Z$), $e \equiv E/A$, of an **uniform system of neutrons (N) and protons (Z)** as a function of the **neutron ($\rho_n = N/V$) and proton ($\rho_p = Z/V$) densities ...**

→ ... **at zero temperature:** room temperature $10^2\text{K} \rightarrow 10^{-8}$ MeV while “cold” neutron stars are at about $10^{10}\text{K} \rightarrow 1$ MeV. **Separation energy** in stable nuclei (equivalent to ionization energy in atoms) is of **several MeV**.

→ ... **unpolarized:** energy favors **couples** of neutrons and protons **occupying the same state but with opposite spins** (equivalent to electrons in atoms)

→ ... **isospin symmetric:** neutron-neutron, proton-proton and neutron-proton **nuclear interaction** are very **similar** among them. **Masses** of neutrons and protons are almost **degenerate**. Hence neutrons and protons can be thought as **two states** of the **same particle** with different isobaric spin or **isospin** (in analogy with spin): the **nucleon**.

→ ... and **no Coulomb:** **idealized uniform system** (focus on strong interaction, much stronger than Coulomb at nuclear scale $\sim 10^{15}$ fm). Real systems are finite and electrically neutral so no divergences in adding Coulomb.

Nuclear Equation of State (EoS)

What is it? How is defined?

Definition: the energy (E) per nucleon ($A=N+Z$), $e \equiv E/A$, of an **uniform system of neutrons (N) and protons (Z)** as a function of the **neutron ($\rho_n = N/V$) and proton ($\rho_p = Z/V$) densities ...**

→ ... **at zero temperature:** room temperature $10^2\text{K} \rightarrow 10^{-8}$ MeV while “cold” neutron stars are at about $10^{10}\text{K} \rightarrow 1$ MeV. **Separation energy** in stable nuclei (equivalent to ionization energy in atoms) is of **several MeV**.

→ ... **unpolarized:** energy favors **couples** of neutrons and protons **occupying the same state but with opposite spins** (equivalent to electrons in atoms)

→ ... **isospin symmetric:** neutron-neutron, proton-proton and neutron-proton **nuclear interaction** are very **similar** among them. **Masses** of neutrons and protons are almost **degenerate**. Hence neutrons and protons can be thought as **two states** of the **same particle** with different isobaric spin or **isospin** (in analogy with spin): the **nucleon**.

→ ... and **no Coulomb:** **idealized uniform system** (focus on strong interaction, much stronger than Coulomb at nuclear scale $\sim 10^{15}$ fm). Real systems are finite and electrically neutral so no divergences in adding Coulomb.

Nuclear Equation of State (EoS)

What is it? How is defined?

Definition: the energy (E) per nucleon ($A=N+Z$), $e \equiv E/A$, of an **uniform system of neutrons (N) and protons (Z)** as a function of the **neutron ($\rho_n = N/V$) and proton ($\rho_p = Z/V$) densities ...**

→ ... **at zero temperature:** room temperature $10^2\text{K} \rightarrow 10^{-8}$ MeV while “cold” neutron stars are at about $10^{10}\text{K} \rightarrow 1$ MeV. **Separation energy** in stable nuclei (equivalent to ionization energy in atoms) is of **several MeV**.

→ ... **unpolarized:** energy favors **couples** of neutrons and protons **occupying the same state but with opposite spins** (equivalent to electrons in atoms)

→ ... **isospin symmetric:** neutron-neutron, proton-proton and neutron-proton **nuclear interaction** are very **similar** among them. **Masses** of neutrons and protons are almost **degenerate**. Hence neutrons and protons can be thought as **two states** of the **same particle** with different isobaric spin or **isospin** (in analogy with spin): the **nucleon**.

→ ... and **no Coulomb:** **idealized uniform system** (focus on strong interaction, much stronger than Coulomb at nuclear scale $\sim 10^{15}$ fm). Real systems are finite and electrically neutral so no divergences in adding Coulomb.

Nuclear Equation of State (EoS)

What is it? How is defined?

Definition: the energy (E) per nucleon ($A=N+Z$), $e \equiv E/A$, of an **uniform system of neutrons (N) and protons (Z)** as a function of the **neutron ($\rho_n = N/V$) and proton ($\rho_p = Z/V$) densities ...**

→ ... **at zero temperature:** room temperature $10^2\text{K} \rightarrow 10^{-8}$ MeV while “cold” neutron stars are at about $10^{10}\text{K} \rightarrow 1$ MeV. **Separation energy** in stable nuclei (equivalent to ionization energy in atoms) is of **several MeV**.

→ ... **unpolarized:** energy favors **couples** of neutrons and protons **occupying the same state but with opposite spins** (equivalent to electrons in atoms)


→ ... **isospin symmetric:** neutron-neutron, proton-proton and neutron-proton **nuclear interaction** are very **similar** among them. **Masses** of neutrons and protons are almost **degenerate**. Hence neutrons and protons can be thought as **two states** of the **same particle** with different isobaric spin or **isospin** (in analogy with spin): the **nucleon**.

→ ... and **no Coulomb: idealized uniform system** (focus on strong interaction, much stronger than Coulomb at nuclear scale $\sim 10^{-15}$ m). Real systems are finite and electrically neutral so no divergences in adding Coulomb.


Nuclear Equation of State (EoS)

Unpolarized, uniform nuclear matter at $T=0$ assuming isospin symmetry


Neutron Star



It is convenient to write the energy per nucleon (e) as a function of the total density $\rho = \rho_n + \rho_p$ and the relative difference $\delta = (\rho_n - \rho_p) / \rho$ for $\delta \rightarrow 0$:




$$e(\rho, \delta) = e(\rho, 0) + S_2(\rho)\delta^2 + S_4(\rho)\delta^4 + \mathcal{O}[\delta^6]$$




Nuclear Equation of State (EoS)

Unpolarized, uniform nuclear matter at T=0 assuming isospin symmetry


Neutron Star



It is convenient to write the energy per nucleon (e) as a function of the total density $\rho = \rho_n + \rho_p$ and the relative difference $\delta = (\rho_n - \rho_p) / \rho$ for $\delta \rightarrow 0$:




$$e(\rho, \delta) = e(\rho, 0) + S_2(\rho)\delta^2 + S_4(\rho)\delta^4 + \mathcal{O}[\delta^6]$$



Nuclear Equation of State (EoS)

Unpolarized, uniform nuclear matter at T=0 assuming isospin symmetry

$$e(\rho, \delta) = e(\rho, 0) + S_2(\rho)\delta^2$$



It is customary to **Taylor expand** $e(\rho, 0)$ and $S(\rho)$ around **nuclear saturation density** $\rho_0 \sim 0.16 \text{ fm}^{-3}$

$$e(\rho, 0) = e(\rho_0, 0) + \frac{1}{2}K_0x^2 + \mathcal{O}[\rho^3] \text{ where } x = \frac{\rho - \rho_0}{3\rho_0}$$

$$S(\rho) = J + Lx + \frac{1}{2}K_{\text{sym}}x^2 + \mathcal{O}[\rho^3, \delta^4]$$

K → how **compressible** is matter @ ρ_0

J → **penalty energy** for systems with $\rho_n \neq \rho_p$ @ $\rho_n + \rho_p = \rho_0$

L → **pressure** for systems with $\rho_n \neq \rho_p$ @ ρ_0

$$P = - \left. \frac{\partial E}{\partial V} \right|_{A=\text{const.}} = \frac{1}{3} \rho \delta^2 L(\rho)$$


Electric Dipole Polarizability: introduction

The **electric dipole polarizability** measures the **tendency** of the nuclear **charge distribution** to be **distorted**


$$\alpha = \frac{\text{electric dipole moment}}{\text{external electric field applied}}$$

Microscopically, it relates with the **photo-absorption cross-section**

$$\alpha_D = \frac{\hbar c}{2\pi^2 e^2} \int \frac{\sigma_{\text{abs}}}{\omega^2} d\omega,$$



Measured using **polarized** proton scattering at **very forward angles** (dominated by **E1** and **M1** well separated)



$$\alpha_D(^{208}\text{Pb}) = 19.6 \pm 0.6 \text{ fm}^3$$

$$\Delta r_{\text{np}} = 0.156^{+0.025}_{-0.021} \text{ fm}$$

Electric Dipole Polarizability: theory

Theoretically, the total photo-absorption cross section, can be written as

$$\sigma_{\gamma\text{-abs}} = 4\pi^2\alpha \sum_{\nu} (E_{\nu} - E_0) |\langle \nu | F_{\text{dipole}} | 0 \rangle|^2$$

Dipole operator
subtract CM motion

And, thus,

$$\alpha_D = 2 \sum_{\nu \neq 0} \frac{|\langle \nu | F_{\text{dipole}} | 0 \rangle|^2}{E_{\nu} - E_0} \equiv 2m_{-1} \quad \frac{eN}{A} \sum_{i=1}^Z r_i Y_{1M}(\hat{r}_i) - \frac{eZ}{A} \sum_{i=1}^N r_i Y_{1M}(\hat{r}_i)$$

Considering the G.S. perturbed by an external field λF (with $\lambda \rightarrow 0$):

$\langle \mathcal{H} \rangle = \langle \mathcal{H}_0 + \lambda F_{\text{dipole}} \rangle$; The variation in the expectation energy can be written as:

$$\delta \langle \mathcal{H} \rangle = \lambda^2 \sum_{\nu \neq 0} \frac{|\langle \nu | F | 0 \rangle|^2}{E_{\nu} - E_0} + \mathcal{O}(\lambda^3) = \lambda^2 m_{-1} + \mathcal{O}(\lambda^3)$$

$$m_{-1} = \frac{1}{2} \frac{\partial^2 \langle \mathcal{H} \rangle}{\partial \lambda^2} \Big|_{\lambda=0}$$

Dielectric
Theorem

Electric Dipole Polarizability: simple model & correlations



Applying the **dielectric theorem** to the **Droplet Model** Hamiltonian (first Migdal and latter on Meyer et al. NPA385, 269) one can find

$$\alpha_D \approx \frac{A \langle r^2 \rangle}{12J} \left[1 + \frac{5 \Delta r_{np} + \sqrt{\frac{3}{5} \frac{e^2 Z}{70J}} - \Delta r_{np}^{\text{surface}}}{\langle r^2 \rangle^{1/2} (I - I_C)} \right]$$

$J = e(\text{PNM}) - e(\text{SNM}) \rightarrow$ Symmetry energy at ρ_0


$\Delta r_{np} \equiv \langle r_n^2 \rangle^{1/2} - \langle r_p^2 \rangle^{1/2} \rightarrow$ Neutron skin thickness

Using microscopic calculations (Energy Density Functionals & Ab initio)



$$\alpha_D J = (301 \pm 32) + (1922 \pm 73) \Delta r_{np}$$

Electric Dipole Polarizability in ^{48}Ca and ^{208}Pb



Combined Theoretical Analysis of the Parity-Violating Asymmetry for ^{48}Ca and ^{208}Pb

Paul-Gerhard Reinhard, Xavier Roca-Maza, and Witold Nazarewicz
Phys. Rev. Lett. **129**, 232501 – Published 2 December 2022


| Observable | z | σ_{exp} | σ_{model} | σ_{method} | σ_{em} | PPD |
|-----------------------------|--------|-----------------------|-------------------------|--------------------------|----------------------|-------------------------|
| $E/A(^{208}\text{Pb})$ | -7.867 | 0 | 0.54 | 0.5 | — | $-8.06^{+0.99}_{-0.88}$ |
| $R_p(^{208}\text{Pb})$ | 5.45 | 0 | 0.17 | 0.05 | — | $5.43^{+0.21}_{-0.23}$ |
| $\alpha_D(^{48}\text{Ca})$ | 2.07 | 0.22 | 0.06 | 0.1 | — | $2.30^{+0.31}_{-0.26}$ |
| $\alpha_D(^{208}\text{Pb})$ | 20.1 | 0.6 | 0.59 | 0.8 | — | $22.6^{+2.1}_{-1.8}$ |

Ab initio predictions link the neutron skin of ^{208}Pb to nuclear forces

Baishan Hu, Weiguang Jiang, Takayuki Miyagi, Zhonghao Sun, Andreas Ekström, Christian Forssén, Gaute Hagen, Jason D. Holt, Thomas Papenbrock, S. Ragnar Stroberg & Ian Vernon

Nature Physics **18**, 1196–1200 (2022) | [Cite this article](#)

Electric Dipole Polarizability: How nuclear models perform for other nuclei?



Parity Violating Asymmetry: introduction


Elastic electron scattering by nuclei:

→ **Parity conserving**

Exchange of γ that essentially couples to protons $Q_p^C=1; Q_n^C=0$

→ **Parity violating**

Exchange Z_0 that essentially couples to neutrons $Q_p^W=0.07; Q_n^W=-0.99$



Ultra-relativistic electrons ($m_e \rightarrow 0$) with spin **aligned** (+) or **anti-aligned** (-) to the beam line, in approaching the nucleus:

$$\left[\vec{\alpha} \cdot \vec{p} + V_{\text{Coulomb}}(\vec{r}) \pm V_{\text{Weak}}(\vec{r}) \right] \Psi_{\pm} = E \Psi_{\pm}$$

$$\sigma \approx \left| \begin{array}{c} \text{Diagram 1: } e^- \text{ and } 208\text{Pb} \text{ connected by a } \gamma \text{ line} \\ + \\ \text{Diagram 2: } e^- \text{ and } 208\text{Pb} \text{ connected by a } Z^0 \text{ line} \end{array} \right|^2$$

One can get **advantage** from this **interference pattern** between **electromag.** and **weak** interact.

$$A_{PV} = \frac{\frac{d\sigma_+}{d\Omega} - \frac{d\sigma_-}{d\Omega}}{\frac{d\sigma_+}{d\Omega} + \frac{d\sigma_-}{d\Omega}} \approx \frac{\text{Weak}}{\text{Coulomb}}$$


Parity Violating Asymmetry: experiment

Lead and Calcium

Radius Experiments

PREx & CREx @ JLab

$$A_{PV} \sim \frac{G_F}{4\sqrt{2}\pi\alpha} Q^2 \sim 10^{-4} (Q[\text{GeV}])^2$$



R. Michaels

→ Spectrometers $\sim 5^\circ$

→ e^- beam $E \sim \text{GeV}$

$$Q_{\text{PREx}}^2 = 0.00616 \text{ GeV}^2$$

$$Q_{\text{CREx}}^2 = 0.0297 \text{ GeV}^2$$

→ Demanding experiment:

For 10^6 electrons or more only **one** of them **interact weakly** with the nucleus

New experiment on ^{208}Pb @ Mainz !!! (MREX)

Parity Violating Asymmetry: theory

To solve the **Dirac equation**


$$\left[\vec{\alpha} \cdot \vec{p} + V_{\text{Coulomb}}(\vec{r}) \pm V_{\text{Weak}}(\vec{r}) \right] \Psi_{\pm} = E \Psi_{\pm}$$

The main inputs are the **electromagnetic** ρ_{ch} and **weak charge** ρ_{W} distributions

$$V_{\text{Coulomb}}(\vec{r}) = Z\alpha \int \frac{\rho_{\text{ch}}(\vec{r}')}{|\vec{r} - \vec{r}'|} d\vec{r}'$$

$$V_{\text{Weak}}(\vec{r}) = \frac{G_F}{2\sqrt{2}} \rho_{\text{W}}(\vec{r})$$

We need to know the **neutron and proton distributions** in nuclei and the **electromagnetic and weak charge form factors** of the **neutron** and the **proton**



Parity Violating Asimmetry: simple model & correlations

The parity violating asymmetry within the PWBA:


$$A_{PV}^{PWBA} = \frac{G_F Q^2}{4\sqrt{2}\pi\alpha} \left[\underbrace{4 \sin^2 \theta_W}_{\approx 1} + \frac{F_n(Q) - F_p(Q)}{F_p(Q)} \right]$$

$$\xrightarrow{Q \rightarrow 0} \frac{G_F Q^2}{4\sqrt{2}\pi\alpha} \left[1 - \frac{Q^2 \langle r_p^2 \rangle}{3} \frac{\Delta r_{np}}{\langle r_p^2 \rangle^{1/2}} \right]$$

30% due to Coulomb distortions (PREX) does not blur the main physics

The parity violating asymmetry within the DWBA:


- ◆ → Energy Density Functionals
- → Expeirmental ρ_n form hadronic probes + ρ_p from e^- scattering



Parity Violating Asymmetry: measurements vs. theory


Selecting those models (we consider) well calibrated to masses and radii

Theoretical (EDFs and *ab initio*) and experimental 1σ errors overlap in ^{208}Pb but not in ^{48}Ca → **No simultaneous description**



Fitting protocol also including A_{PV} and α_{D}

- SV-min
- SV-APV $^2\alpha^2$
- SV-APV $^1\alpha^2$
- SV- α^2
- SV-min*
- RD-min
- RD-APV $^2\alpha^2$
- RD-APV $^1\alpha^2$
- RD- α^2
- PC-min




Combined Theoretical Analysis of the Parity-Violating Asymmetry for ^{48}Ca and ^{208}Pb

Paul-Gerhard Reinhard, Xavier Roca-Maza, and Witold Nazarewicz
Phys. Rev. Lett. **129**, 232501 – Published 2 December 2022

Magenta dashed lines from extrapolated Δr_{np} given in G. Hagen et al. Nature Physics 12, 186–190 (2016) and H. Bu et al. Nature Physics (2022)

Summary: model performance

A_{PV} (sensitive to Δr_{np}) and α_D (sensitive to J and Δr_{np}) in ^{48}Ca and ^{208}Pb




Simultaneous description of dipole polarizabilities → point to a **good understanding of symmetry energy (J) and neutron skins (Δr_{np})**

Ab-initio (B. Hu) Nature Physics (2022)

$$\alpha_D(^{48}\text{Ca}) \quad 2.30^{+0.31}_{-0.26}$$

$$\alpha_D(^{208}\text{Pb}) \quad 22.6^{+2.1}_{-1.8}$$

No simultaneous description of parity violating asymmetries (ground state observable) → point to a **deficient understanding of neutron skins**



Magenta dashed lines from extrapolated Δr_{np} given in G. Hagen et al. Nature Physics 12, 186–190 (2016) and H. Bu et al. Nature Physics (2022)

A_{pV} theoretical corrections?

In all analysis: Coulomb potential has been considered at tree level while Q_{Weak} has been corrected at one-loop level (interference of the γ and Z_0 exchange):

$$[-i\vec{\alpha} \cdot \vec{\nabla} + V_{\text{Coul}} \pm V_{\text{Weak}}]\Psi_{\pm} = E\Psi_{\pm}$$

First order QED includes: vacuum polarization (VP), self-energy and e^- vertex correction (V-SE) [Jakubassa-Amundsen (2024) JPG: Nucl. Part. Phys. 51 035105]

$$[-i\vec{\alpha} \cdot \vec{\nabla} + V_{\text{Coul}} \pm V_{\text{Weak}} + V_{\text{VP}} + V_{\text{V-SE}}]\Psi_{\pm} = E\Psi_{\pm}$$


Dispersion corrections: interference of the γ and Z_0 exchange have been shown to be relevant for estimating the Q_{weak} .

On A_{pV} @ PREx kin., corrections **small & cancel**

- Box diagram -0.1%
- QED 0.09%

[Jakubassa-Amundsen and XRM, arXiv:2507.15380]


[XRM and Jakubassa-Amundsen PRL 134, 192501 (2025)]



Isobaric Analog State: Example + theory

$$F = T_{\pm} = \sum_i^A t_{\pm}(i)$$

Charge-exchange experiments



Isospin algebra analogous to spin algebra $s \rightarrow t$ and $\tau \rightarrow \sigma$ (Pauli matrices with $t = \tau/2$)

$$t_- |n\rangle = \frac{1}{2} |p\rangle$$

$$t_+ |p\rangle = -\frac{1}{2} |n\rangle$$

$$T_+^\dagger = T_- \quad T_-^\dagger = T_+$$

$$[T_z, T_{\pm}] = \pm T_{\pm}$$

$$[T_+, T_-] = 2T_z$$

→ non-energy weighted sum rule:

$$\begin{aligned}
 m_0^- - m_0^+ &= \langle 0 | T_+ T_- | 0 \rangle - \langle 0 | T_- T_+ | 0 \rangle \\
 &= \langle 0 | [T_+, T_-] | 0 \rangle = \langle 0 | 2T_z | 0 \rangle \\
 &= N - Z
 \end{aligned}$$

Note: If not isospin-mixing it would be zero!!

→ energy weighted sum rule:

$$m_1 = \sum_{\nu} (E_{\nu} - E_0) |\langle \nu | F | 0 \rangle|^2 = \langle 0 | T_+ [\mathcal{H}, T_-] | 0 \rangle$$

$[\mathcal{H}, T_-] \neq 0$ only if \mathcal{H} contains terms that **breaks isospin invariance**

Isobaric Analog State: Theory

$$F = T_{\pm} = \sum_i^A t_{\pm}(i)$$

→ Hence, the **centroid energy** m_1/m_0 :

$$E_{\text{IAS}} = \frac{\langle 0 | T_+ [\mathcal{H}, T_-] | 0 \rangle}{\langle 0 | T_+ T_- | 0 \rangle} = \frac{1}{N - Z} \langle 0 | T_+ [\mathcal{H}, T_-] | 0 \rangle$$

→ Assuming a simple model: **independent particle** model with only **Coulomb breaking isospin symmetry** (neglect exchange effects)

$$E_{\text{IAS}}^{\text{C,direct}} = \frac{1}{N - Z} \int [\rho_n(\vec{r}) - \rho_p(\vec{r})] U_{\text{C}}^{\text{direct}}(\vec{r}) d\vec{r},$$

$$U_{\text{C}}^{\text{direct}}(\vec{r}) = \int \frac{e^2}{|\vec{r}_1 - \vec{r}|} \rho_{\text{ch}}(\vec{r}_1) d\vec{r}_1$$


→ Assuming sharp sphere to describe ρ_n and ρ_p and $\rho_{\text{ch}} = \rho_p$

$$U_{\text{C}}^{\text{direct}}(\vec{r}) = \begin{cases} \frac{Ze^2}{2R_p} \left(3 - \frac{r^2}{R_p^2} \right) & \text{for } r < R_p \\ \frac{Ze^2}{r} & \text{for } r > R_p \end{cases}$$

$$\begin{aligned} E_{\text{IAS}} &\approx E_{\text{IAS}}^{\text{C,direct}} \\ &\approx \frac{6Ze^2}{5R_p} \left(1 - \frac{1}{2} \frac{N}{N-Z} \frac{R_n - R_p}{R_p} \right) \\ &\approx \frac{6}{5} \frac{Ze^2}{r_0 A^{1/3}} \left(1 - \sqrt{\frac{5}{12}} \frac{N}{N-Z} \frac{\Delta R_{\text{np}}}{r_0 A^{1/3}} \right) \end{aligned}$$

Isobaric Analog State:

Theory+experiment



Exp errors in IAS ~
tens of keV


(or smaller)

Exp width IAS

~ hundreds of keV

SAMi-ISB: includes ISB effects due to Coulomb exchange, QED corrections and nuclear ISB

Summary on Δr_{np} in ^{208}Pb :



Summary



Different ways to learn about the **neutron distribution** in atomic nuclei using **different observables** provide **different answers**

However, within **1.5–2 σ** **all looks fine**, both from phenomenologic **Energy Density Functionals** and from **Ab Initio**

From Theory:

- An effort to better understand the **parity violating asymmetry** and the **beam normal spin asymmetry**^[*] in **^{208}Pb** is needed [PREX & CREX, PRL 128, 142501 (2022)].
- An effort to better understand the **systematics** on the **dipole polarizability** would be of interest [see e.g. Sn chain in Bassauer et al. PLB 810, 135804 (2020)].
- An effort to **understand ISB effects in the medium** from first principles [see e.g. Sagawa et al. Phys. Rev. C 109, L011302 (2024)]

From Experiment (low-energy):



- An effort to improve the accuracy in the **parity violating asymmetry in ^{208}Pb** (and measure **other Q values**) **is needed**. Measuring other nuclei would also be desirable.
- **Systematic** measurements of the **dipole polarizability** along **neutron rich** isotopic chains (e.g. $N > 74$ Sn isotopes) could help testing models and improve our understanding of this observable.

Collaborators

- Gianluca **Colò** (University of Milan)
- Pietro **Klausner** (University of Milan)
- Tomoya **Naito** (University of Tokyo)
- Xavier **Vinyes** & Mario **Centelles** (University of Barcelona)
- Jorge **Piekarewicz** (Florida State University)
- Nils **Paar** & Dario **Vretenar** (University of Zagreb)
- Bijay K. **Agrawal** (Saha Institute of Nuclear Physics)
- P.-G. **Reinhard** (University of Erlangen-Nürnberg)
- Witold **Nazarewicz** (FRIB and Michigan State University)
- Doris H. **Jakubassa-Amunsden** (Ludwig-Maximilians-Universität München)

RCNP - proton elastic scattering

Polarized proton elastic scattering @ 295 MeV sensitive to the overall strong size of the nucleus



$$\Delta r_{\text{np}} = 0.211^{+0.054}_{-0.063} \text{ fm}$$

Neutron density distributions of $^{204,206,208}\text{Pb}$ deduced via proton elastic scattering at $E_p = 295$ MeV


J. Zenhiro^{1*}, H. Sakaguchi^{1,2}, T. Murakami¹, M. Yosoi^{1,2}, Y. Yasuda^{1,2}, S. Terashima^{1,4}, Y. Iwao¹, H. Takeda², M. Itoh^{3,5} et al.

Show more

Phys. Rev. C 82, 044611 - Published 22 October, 2010

LHC - Relativistic Heavy Ion Collisions

Particle distributions and **collective flow** in $^{208}\text{Pb} + ^{208}\text{Pb}$ RHIC @ LHC are **sensitive** to the overall strong **size** of the nuclei involved.



Determination of the Neutron Skin of ^{208}Pb from Ultrarelativistic Nuclear Collisions



Giuliano Giacalone¹, Govert Nijss², and Wilke van der Schee^{3,4}

Show more

Phys. Rev. Lett. 131, 202302 - Published 15 November, 2023

$$\Delta r_{np} = 0.217 \pm 0.058 \text{ fm}$$

Summary on Δr_{np} in ^{48}Ca and ^{208}Pb



R_{ch}^4 inferred from expt. R_{ch}^2
 R_{ch}^4 inferred from theor. R_{ch}^2


$\Delta\text{NNLO}_{\text{GO}}$
 1.8/2.0 (EM7.5)

Nonimplausible Hamiltonians

Ab initio computations of the fourth-order charge density moments of ^{48}Ca and ^{208}Pb

Beam normal spin asymmetry (A_n) (aka Alaying power or Sherman function)

Is the asymmetry in the scattering cross section when the incident electron beam is polarized perpendicular (longitudinal $\rightarrow A_{pv}$) to the scattering plane



\Rightarrow Important to understand A_{pv} since it represents a potentially large systematic correction to measured asymmetries.

In Born approximation (one-photon exchange), $A_n = 0$ because time-reversal symmetry forbids such an asymmetry.

Arises from interference between one-photon exchange and the absorptive part of two-photon exchange amplitudes.

New Measurements of the Transverse Beam Asymmetry for Elastic Electron Scattering from Selected Nuclei

[S. Abrahamyan](#)⁴⁵, [A. Acha](#)¹⁰, [A. Afanasev](#)¹¹, [Z. Ahmed](#)³³, [H. Albataineh](#)⁶, [K. Aniol](#)³, [D. S. Armstrong](#)⁷, [W. Armstrong](#)³⁵, [J. Arrington](#)¹ et al. (HAPPEX and PREX Collaborations)

Show more

Phys. Rev. Lett. **109**, 192501 – Published 5 November, 2012
DOI: <https://doi.org/10.1103/PhysRevLett.109.192501>

New Measurements of the Beam-Normal Single Spin Asymmetry in Elastic Electron Scattering over a Range of Spin-0 Nuclei

[D. Adhikari](#)¹, [H. Albataineh](#)², [D. Androic](#)³, [K. Aniol](#)⁴, [D. S. Armstrong](#)⁵, [T. Averett](#)⁵, [C. Ayerbe Gayoso](#)⁵, [S. Barcus](#)⁵, [V. Bellini](#)⁷ et al. (PREX and CREX Collaborations)


Show more

Phys. Rev. Lett. **128**, 142501 – Published 8 April, 2022
DOI: <https://doi.org/10.1103/PhysRevLett.128.142501>



Parity Violating Asimmetry: theory

The main uncertainties come from nuclear model errors on the description of the neutron distribution (blue error bars)

→ Coulomb distortions are important (order αZ)



Are γZ_0 box corrections important? not included




Hadronic uncertainties less relevant

| Parameter | Value |
|--|----------------------|
| $\langle r_p^2 \rangle$ (fm ²) | 0.726 ± 0.019 |
| $\langle r_n^2 \rangle$ (fm ²) | -0.1161 ± 0.0022 |
| μ_p | 2.792 847 |
| μ_n | -1.9130 |
| $Q_p^{(W)}$ | 0.0713 ± 0.0001 |
| $Q_n^{(W)}$ | -0.9888 ± 0.0011 |
| ρ_s | -0.24 ± 0.70 |
| κ_s | -0.017 ± 0.004 |
| $Q_{126,82}^{(W)}$ | -117.9 ± 0.3 |

Bayesian inference Skyrme EDF: can we accommodate α_D and A_{PV} without compromising other observables?


By P. Klausner



Bayesian inference Skyrme EDF: can we accommodate α_D and A_{PV} without compromising other observables?

Pietro Klausner, Gianluca Colò, Xavier Roca-Maza, and Enrico Vigezzi
Phys. Rev. C 111 014311 (2025)

By P. Klausner




| | B.E. (MeV) | R_{ch} (fm) | ΔE_{SO} (MeV) |
|-------------------|----------------|-----------------|-----------------------|
| ^{208}Pb | 1636 ± 1.8 | 5.49 ± 0.03 | 2.34 ± 0.16 |
| ^{48}Ca | 417 ± 1.2 | 3.51 ± 0.02 | 1.92 ± 0.20 |
| ^{40}Ca | 342 ± 1.6 | 3.50 ± 0.02 | — |
| ^{56}Ni | 482 ± 1.4 | — | — |
| ^{68}Ni | 590 ± 1.0 | — | — |
| ^{100}Sn | 826 ± 1.6 | — | — |
| ^{132}Sn | 1103 ± 1.7 | 4.71 ± 0.03 | — |
| ^{90}Zr | 784 ± 1.3 | 4.27 ± 0.02 | — |

Isoscalar resonances

| | E_{GMR}^{IS} (MeV) | E_{GQR}^{IS} (MeV) |
|-------------------|----------------------|----------------------|
| ^{208}Pb | 13.5 ± 0.3 | 10.8 ± 0.4 |
| ^{90}Zr | 17.8 ± 0.4 | — |

| | α_D (fm ³) | $m(1)$ (MeV fm ²) | A_{PV} (ppb) |
|-------------------|-------------------------------|-------------------------------|----------------|
| ^{208}Pb | 19.5 ± 0.5 | 958 ± 22 | 589 ± 5 |
| ^{48}Ca | 2.30 ± 0.08 | — | 2591 ± 54 |



Keeping ground and excited state properties within typical Skyrme-EDF accuracy